

Vaejoidea Species	LAS: Right/Left	Vaejoidea Species	LAS: Right/Left
<i>Vejoidea, Smeringurus, Paruroctonus, Paravaejoidea</i>		<i>Vaejoidea</i> “eusthenura” group and <i>Syntropis</i>	
<i>Vejoidea longiunguis</i> * (S)	9–21 (11.833) [24]	<i>Vaejoidea coahuilae</i>	4/6
<i>Smeringurus aridus</i> (S)	traces of 2	<i>V. confusus</i> (S)	9/7
<i>S. grandis</i>	1/1, 2/2, 3/3, 3/3	<i>V. diazi</i> (S)	6/5, 7/6, x_6
<i>S. mesaensis</i> * (S)	4–10 (7.406) [32]	<i>V. eusthenura</i> (S)	8/7, 5/6
<i>S. v. immanis</i>	4/x	<i>V. globosus</i> (S)	6/4, 2/3
<i>Paruroctonus arnaudi</i> (S)	9/11, 12/11, 13/12	<i>V. gravicaudus</i> (S)	8/9, 3/4
<i>P. arenicola nupides</i>	6/6	<i>V. hoffmanni</i>	6/5, 6/7, 4/7
<i>P. bantai saratoga</i>	2/2, 3/3	<i>V. punctatus</i> * (S)	2–8 (4.727) [33]
<i>P. becki</i> (S)	5/5, 8/8	<i>V. puritanus</i> * (S)	3–12 (7.714) [28]
<i>P. boreus</i> (S)	4/4, 7/x	<i>V. spinigerus</i> (S)	4/x, 5/3, 7/5, x/3
<i>P. borregoensis</i>	7/8	<i>V. viscaimensis</i> * (S)	6–15 (9.385) [26]
<i>P. gracilior</i> * (S)	4–9 (6.240) [25]	<i>V. vittatus</i> (S)	7/x
<i>P. hirsutipes</i>	ABSENT	<i>V. waeringi</i> (S)	6/11, 5/7
<i>P. luteolus</i> (S)	7/6, 6/7	<i>Syntropis macrura</i>	5/6, 10/8, 6/7, 3/3
<i>P. silvestrii</i> * (S)	1–10 (6.643) [28]	<i>Franckeus</i> and <i>Vaejoidea</i> “nigrescens” group	
<i>P. stahnkei</i>	6/7	<i>Franckeus minckley</i>	9/8, 7/5, 6/5
<i>P. surensis</i> (S)	9/8, 11/10, 8/12	<i>F. pennisularis</i>	8/7, 3/4, 5/6, 3/3
<i>P. ventosus</i> (S)	14/10, 10/11	<i>Vaejoidea davidi</i>	ABSENT
<i>P. utahensis</i> (S)	5/6, 3/4	<i>V. decipiens</i>	ABSENT
<i>P. xanthus</i>	1/1	<i>V. janssi</i>	x/2, 3/5, 5/4, 7/6, 7/9, 2/4
<i>Paravaejoidea pumilis</i> * (S)	1–7 (4.306) [36]	<i>V. nigrescens</i>	traces of 2
<i>Stahnkeus</i> and <i>Serradigitus</i>		<i>V. pococki</i>	ABSENT
<i>Stahnkeus deserticola</i>	4/x	<i>V. solegladi</i>	ABSENT
<i>S. harbisoni</i>	2/5	<i>Vaejoidea</i> “mexicanus” group	
<i>S. subtilimanus</i>	1/1, 4/3	<i>Vaejoidea carolinianus</i>	2/x
<i>Serradigitus adcocki</i>	2/1	<i>V. granulatus</i> (S)	3/2
<i>S. baueri</i>	5/6	<i>V. jonesi</i>	5/5, 4/5
<i>S. calidus</i>	5/4	<i>V. lapidicola</i>	3/3
<i>S. gertschi</i> (S)	7/6, 10/11	<i>V. mexicanus</i>	ABSENT
<i>S. joshuaensis</i> * (S)	3–8 (5.409) [22]	<i>V. paysonensis</i>	7/5, 8/7, 4/5
<i>S. littoralis</i> (S)	3/3, 4/4	<i>V. vorhiesi</i> *	1–7 (5.000) [14]
<i>S. minutis</i>	6/7	<i>Pseudouroctonus</i> and <i>Uroctonites</i>	
<i>S. torridus</i>	4/4, 5/5	<i>Pseudouroctonus andreas</i> * (S)	2–5 (3.643) [14]
<i>S. wupatkiensis</i>	5/5	<i>P. angelenus</i>	2/x
<i>Vaejoidea</i> “punctipalpi” group		<i>P. apacheanus</i> (S)	ABSENT
<i>V. bruneus</i>	6/7	<i>P. iviei</i>	ABSENT
<i>V. cazieri</i> (S)	4/4, 4/5	<i>P. minimus castaneus</i>	traces x/3
<i>V. hirsuticauda</i> (S)	7/9, 8/x	<i>P. minimus thompsoni</i>	3/2, 4/3, 3/3
<i>V. intrepidus</i>	4/7, 4/5	<i>P. reddelli</i>	3/x
<i>V. magdalensis</i> * (S)	4–8 (5.357) [14]	<i>Uroctonites guiliani</i>	ABSENT
<i>V. occidentalis</i>	ABSENT	<i>U. huachuca</i> (S)	ABSENT
<i>V. punctipalpi</i>	4/5	<i>U. monterius</i>	ABSENT
<i>V. russelli</i>	8/10		

Table 1: Laterobasal Aculear Serrations (LAS) statistics for family Vaejoidea showing number of denticles per lateral surface. Although numbers were determined from ventral view of telson, right/left delineation is from a dorsal perspective. * Indicates species where larger number of specimens were examined thus a min/max (mean) [number of samples] is provided (see Table 2 for additional data). Note, each side of the aculeus is considered a sample. (S) = LAS presence/absence verified with SEM.

quite minute in size; individual denticles are between 17 and 36 micrometers in length. An individual denticle is shaped as a rounded tooth basally narrowing distally into a point, angled approximately 35 degrees from the surface of the aculeus, pointing towards the aculeus tip (Figs. 30–33). The longest line of denticles, found in *Vejoidea longiunguis* (Figs. 2–5) whose aculeus is quite long and slender, was less than 0.5 mm in length and consisted of 14 serrated denticles. Since the LAS is positioned in the area where the aculeus starts darkening, we see that the base of the denticle row is darkened as well, giving a somewhat overall pigmented look to the LAS.

The composition and spacing of individual denticles is quite variable. There are cases where the individual denticles are densely organized basally and then the spacing becomes larger at the distal aspect, sometimes the most distal denticle being quite separated from the previous ones (Fig. 22). Individual denticles are sometimes smaller basally, or they can be irregularly doubled as seen in Fig. 33 for *Paruroctonus ventosus*.

It must be stressed here that the LAS structure does not necessarily occur at the same relative position on the aculeus. Instead, its location is consistently at the region where the narrowing and darkening of the aculeus occurs, and is always distal of the VDSP and DDSP,

Species/Genders	Number of Denticles in LAS
<i>Paravaejovis pumilis</i>	1-7 (4.306) (± 1.238) [36] {3.068-5.544} \rightarrow 0.288
Males	2-7 (4.500)
Females	1-4 (3.333)
<i>Vejovoidus longiunguis</i>	9-21 (11.833) (± 3.031) [24] {8.802-14.865} \rightarrow 0.256
Males	10-21 (13.500)
Females	9-16 (11.000)
<i>Smeringurus mesaensis</i>	4-10 (7.406) (± 1.720) [32] {5.686-9.126} \rightarrow 0.232
Males	5-10 (7.944)
Females	4-9 (6.875)
<i>Paruroctonus gracilior</i>	4-9 (6.240) (± 1.332) [25] {4.908-7.572} \rightarrow 0.213
Males	4-9 (6.261)
Females	6 (6.000)
<i>Paruroctonus silvestrii</i>	1-10 (6.643) (± 2.264) [28] {4.379-8.907} \rightarrow 0.341
Males	1-10 (6.350)
Females	5-8 (7.375)
<i>Vaejovis viscaïnensis</i>	6-15 (9.385) (± 2.210) [26] {7.174-11.595} \rightarrow 0.236
Males	6-15 (9.944)
Females	6-12 (8.125)
<i>Vaejovis punctatus</i>	2-8 (4.727) (± 1.506) [33] {3.222-6.233} \rightarrow 0.319
Males	2-8 (5.190)
Females	2-6 (3.917)
<i>Vaejovis puritanus</i>	3-12 (7.714) (± 1.630) [28] {6.085-9.344} \rightarrow 0.211
Males	3-12 (7.579)
Females	9-10 (9.500)
<i>Vaejovis magdalensis</i>	4-8 (5.357) (± 1.277) [14] {4.080-6.635} \rightarrow 0.238
Males	4-8 (5.357) [14]
Females	-
<i>Serradigitus joshuaensis</i>	3-8 (5.409) (± 1.368) [22] {4.041-6.777} \rightarrow 0.253
Males	-
Females	3-8 (5.409)
<i>Vaejovis vorhiesi</i>	1-7 (5.000) (± 1.569) [14] {3.431-6.569} \rightarrow 0.314
Males	5-7 (5.500)
Females	1-7 (4.800)
<i>Pseudouroctonus andreas</i>	2-5 (3.643) (± 0.745) [14] {2.898-4.388} \rightarrow 0.204
Males	2-5 (3.583)
Females	4 (4.000)

Table 2: Statistical data of the number of denticles in the Laterobasal Aculear Serrations (LAS) structure for select species of family Vaejoidea spanning major genera and *Vaejovis* groups. Statistical data group = minimum-maximum (mean) (\pm SDEV) [N] {mean-SDEV – mean+SDEV} \rightarrow coefficient of variability (i.e., SDEV/mean).

whose location is also dependent on the aculeus narrowing. Therefore, for those species with long thin aculeus with little curving (e.g., *Vejovoidus longiunguis*), the LAS occurs more “mid-stinger” (Figs. 2–5). However, on short highly curved stinger scorpions (e.g., *Pseudouroctonus andreas*), the LAS is located towards the distal tip, just where the aculeus abruptly curves.

LAS statistical analysis

The LAS is found on both males and females as well as on juveniles, even instar-2 individuals. There is no indication that the number of denticles per lateral side increases as a scorpion matures. For example, in *Smeringurus mesaensis*, the largest denticle number (for

one side of the LAS) encountered was ten, found both on adults and instar-2 juveniles. In some groups, as indicated elsewhere, we noticed that the LAS structure was more prevalent in juveniles, sometimes not visible in the limited number of adults examined. Although one detects gross trends in the number of denticles across the various vaejoideid groups, the number of denticles is quite variable within a species even specimens from the same locality (see Tables 1 and 2). The number of denticles per LAS side exhibits asymmetry in many cases, the largest mismatch encountered was five in *Vaejovis viscaïnensis* (LAS denticle numbers 7/12).

Table 2 presents statistical data on the number of denticles in the LAS for twelve vaejoideid species. From these data, it is immediately apparent that there is great variability in the number of denticles within a species,

reflecting relatively large standard deviations and coefficients of variability. In many of the species we see that males exhibit on an average higher LAS denticle numbers than females, showing a 14.6 to 35.0 % higher number (based on comparison of the means).

The largest LAS denticle number was found in a male *Vejovoidus longiunguis* exhibiting 19/21, with the species averaging 11.833 [24]. *Vaejovis viscaianensis* exhibited the next highest number, a male with 15/12, and an average of 9.385 [26]. Although these two species are medium to large in size, we also see somewhat high LAS denticle numbers in the smaller species: *Paruroctonus surensis*, males 11/10 and 8/12, and *P. ventosus*, a male with 14/10. As expected, however, some of the smaller species in Vaejoidea do exhibit somewhat small LAS denticle numbers, *Pseudouroctonus andreas* with an average of 3.643 [14], *Serradigitus joshuaensis* with numbers ranging 3 to 8, and *S. littoralis* with 3 to 4.

We calculated the average length of an individual LAS denticle by dividing the length of the LAS row by the number of denticles found in that row (note, this length includes the spacing between denticles which can be variable). Species with the largest average individual denticle are *Smeringurus mesaensis* (35.5 micrometers) and *Vejovoidus longiunguis* (32.6), and species with the smallest average denticle are *Pseudouroctonus andreas* (17.0) and *Paruroctonus ventosus* (20.8). Across the family, the denticle row length ranged 68–456 (184.67) [30] micrometers and the length of an individual denticle (including spacing) ranged 17–35.5 (25.98) [30] micrometers.

LAS trends within Vaejoidea

Table 1 presents a complete synopsis of LAS denticle numbers for all vaejoidean species analyzed in this study. In many cases, multiple specimens were examined, and for those cases where larger numbers of specimens were examined, we show the statistical data group as stated in Table 2. The species in Table 1 are organized in established and/or presumed related genera and/or *Vaejovis* groups.

For genera *Vejovoidus*, *Smeringurus*, *Paruroctonus*, and *Paravaejovis* (Figs. 2-13, 30-31, 33), we detected the LAS structure in all examined species except for *Paruroctonus hirsutipes* (the latter based on a single specimen), in total 21 species. Genus *Vejovoidus* had the largest number of LAS denticles among all species examined, with a standard deviation range of 9–15 (Figs. 2–5, 7). Interestingly, the LAS development was somewhat reduced in the large species genus *Smeringurus*, with only *S. mesaensis* exhibiting somewhat average to high numbers, with a standard deviation range of 6–9 (Fig. 9). Other *Smeringurus* species, especially *S. aridus* and *S. vachoni*, showed

highly reduced LAS structures in some specimens, and it was absent in others. *S. grandis* consistently exhibited LAS denticles but in small numbers, 1–3. Species of *Paruroctonus*, especially where multiple specimens were available, had a well developed LAS structure with numerous denticles (Figs. 8, 10–13). The lithophiles *P. arnaudi*, *P. ventosus*, and *P. surensis* had somewhat high numbers of 9–13, 10–14, and 8–12, respectively. For two species of *Paruroctonus*, we calculated statistical data from multiple specimens providing standard deviation ranges of 5–8 for *P. gracilior* and 4–9 for *P. silvestrii*. The small monotypic genus *Paravaejovis* also exhibited a well developed LAS structure (Fig. 6) with a standard deviation range of 3–6.

All species examined in tribe Stahnkeini (12 species) had the LAS structure (Figs. 1, 18–19, 27, 32) spanning a somewhat moderate denticle number ranging 1–11, the highest number being found in *Serradigitus gertschi striatus*. We provide statistical data on a series of specimens of *S. joshuaensis* (Tab. 2), one of the smallest species in Vaejoidea, showing a standard deviation range of 4–7. Interestingly, the larger species of the genus *Stahnkeus* and of *Serradigitus adcocki* had somewhat small LAS denticle numbers, ranging from 1 to 5 in the material examined.

We examined eight species in the *Vaejovis* “punctipalpi” group where the LAS structure was found in all species except for *V. occidentalis* (based on the examination of a single adult) (Figs. 20–21, 29). The LAS structure development (determined by its number of denticles) is a little more pronounced in this *Vaejovis* group than in tribe Stahnkeini, exhibiting numbers 4–10 (5.677) [31]. For the species *V. magdalensis* we examined several specimens which had a standard deviation range of 4–7.

Fourteen species of the *Vaejovis* “eusthenura” group and genus *Syntropis* were examined, all exhibiting the LAS structure (Figs. 14–17, 22–23, 26, 28). The LAS development of this group of species was the same as that seen in the “punctipalpi” group, with LAS denticle numbers ranging from 2–13. The species *Vaejovis viscaianensis* (Figs. 1, 22) exhibited the largest number of denticles, otherwise; the number of denticles ranged 2–11. Of these species, *V. viscaianensis* is a psammophile, exhibiting some elongation of the pedipalps, metasomal segments and the telson. There seems to be a trend in psammophiles having more developed LAS structures as evidenced in *Vejovoidus*, with the most developed LAS, *Paruroctonus arnaudi* (Fig. 10), and even some of the smaller *Paruroctonus* species, such as *P. ventosus* (Fig. 11) and *P. surensis* (Fig. 12). We calculated statistical data for three species in this group, exhibiting standard deviation ranges of 7–12 for *V. viscaianensis*, 3–6 for *V. punctatus*, and 6–9 for *V. puritanus*. For genus *Syntropis*, a lithophile, we see a similarly developed LAS with denticle numbers ranging 3–10 (6.000) [8].

Pseudochactida: Pseudochactoidea	<i>Opisthophthalmus wahlbergi</i>
Pseudochactidae: <i>Pseudochactas ovchinnikovi</i> (S)	<i>Pandinus imperator</i>
Buthida: Buthoidea	<i>Scorpio maurus</i> (S)
Buthidae: <i>Alayotityus nanus</i>	<i>Urodacus manicatus</i>
<i>Androctonus bicolor</i>	Hemiscorpiidae: <i>Cheloctonus</i> sp.
<i>Anomalobuthus rickmersi</i>	<i>Liocheles australasiae</i> (S)
<i>Buthacus yotvatensis</i>	<i>Liocheles karschii</i>
<i>Buthus occitanus</i>	<i>Hadogenes troglodytes</i>
<i>Centruroides exilicauda</i>	<i>Heteroscorpion goodmani</i>
<i>Centruroides suffusus</i> (S)	<i>Opisthacanthus lepturus</i>
<i>Compsobuthus matthiesseni</i>	Bothriuridae: <i>Bothriurus araguayae</i> (S)
<i>Grosphus hirtus</i>	<i>Bothriurus burmeisteri</i>
<i>Hottentotta minax</i>	<i>Brachistosternus ehrenbergii</i>
<i>Isometrus maculatus</i>	<i>Centromachetes pocockii</i>
<i>Leiurus quinquestriatus</i> (S)	<i>Cercophonius squama</i>
<i>Liobuthus kessleri</i>	<i>Lisposoma josehermana</i>
<i>Lychas</i> sp.	<i>Phoniocercus pictus</i>
<i>Lychas mucronatus</i> (S)	<i>Urophonius paynensis</i>
<i>Mesobuthus caucasicus</i>	Iurida: Chactoidea
<i>Mesobuthus eupeus</i> (S)	Chactidae: <i>Anuroctonus pococki bajae</i> (S)
<i>Microbuthus</i> sp.	<i>Anuroctonus pococki pococki</i>
<i>Microtityus jaumi</i>	<i>Belisarius xambeui</i>
<i>Orthochirus gromovi</i> (S)	<i>Brotheas granulatus</i>
<i>Parabuthus</i> sp.	<i>Broteochactas porosus</i>
<i>Polisius persicus</i>	<i>Chactas exsul</i>
<i>Razianus zarudnyi</i>	<i>Hadrurochactas schaumii</i>
<i>Rhopalurus junceus</i> (S)	<i>Neochactas delicatus</i>
<i>Tityus nematochirus</i>	<i>Nullibrotheas allenii</i> (S)
<i>Uroplectes vittatus</i>	<i>Teuthraustes oculatus</i>
Microcharmidae: <i>Microcharmus hauseri</i>	<i>Uroctonus m. mordax</i> (S)
Chaerilida: Chaeriloidea	<i>Uroctonus m. pluridens</i>
Chaerilidae: <i>Chaerilus celebensis</i> (S)	<i>Vachoniochactas</i> sp.
<i>Chaerilus variegatus</i>	Euscorpiidae: <i>Alloscorpiops lindstroemii</i>
Iurida: Iuroidea	<i>Chactopsis insignis</i>
Caraboctonidae: <i>Caraboctonus keyserlingi</i>	<i>Euscorpiops</i> sp.
<i>Hadruidoidea charcasus</i>	<i>Euscorpius flavicaudis</i>
<i>Hadruidoidea maculatus</i>	<i>Euscorpius gamma</i> (S)
<i>Hadrurus concolorous</i>	<i>Euscorpius italicus</i>
<i>Hadrurus obscurus</i>	<i>Euscorpius mingrelicus</i>
<i>Hoffmannihadrurus aztecus</i>	<i>Euscorpius sicanius</i> (S)
Iuridae: <i>Calchas nordmanni</i>	<i>Megacormus gertschi</i>
<i>Iurus dufourensis</i>	<i>Neoscorpiops tenuicauda</i>
Iurida: Scorpionoidea	<i>Plesiochactas dilutus</i>
Scorpionidae: <i>Bioculus comondae</i> (S)	<i>Scorpiops</i> sp.
<i>Didymocentrus leseurii</i>	<i>Troglocormus willis</i>
<i>Heterometrus longimanus</i>	Superstitioniidae: <i>Superstitionia donensis</i> (S)
<i>Nebo hierichonticus</i>	Vaejovidae: (see Table 1)

Table 3: List of non-vaejovid scorpions where Laterobasal Aculear Serrations (LAS) is *absent*. All parvorders, superfamilies and families are represented (see Table 1 for Vaejovidae). The 73 genera examined are grouped alphabetically. (S) = LAS absence verified with SEM.

Eight species of genus *Franckeus* and the *Vaejovis* “nigrescens” group were examined, among which the two *Franckeus* species and two species of the “nigrescens” group exhibited the LAS structure. In *Franckeus*, we detected well developed LAS, with denticle numbers ranging 3–9 (5.643). For the species *Vaejovis janssi*, only juveniles exhibited the LAS

structure, some with somewhat high numbers of denticles (e.g., 7/6, 7/9). For the other “nigrescens” group species the number of specimens examined was quite limited, some only represented by a solitary specimen, and all were adults. Examination of additional species and material is required for these groups, especially small subadults and juveniles.

In the *Vaejovis* “mexicanus” group (seven species examined, Fig. 25) we encountered the LAS structure in all species, except *V. mexicanus*, which was represented by adults only. For the “vorhiesi” subgroup (i.e., *V. vorhiesi*, *V. paysonensis*, and *V. lapidicola*), all somewhat small species, we see moderately developed LAS, with denticle numbers ranging 1–8 (5.091).

For genera *Pseudouroctonus* and *Uroctonites* (ten species examined) we found the LAS structure in five of the species. *Pseudouroctonus andreas* (Fig. 24) had an average of 3.543 denticles (see Tab. 2) and *P. minimus thompsoni*, 2–4 (found only on juveniles). Species *P. angeleus*, *P. reddelli*, and *P. minimus castaneus* exhibited only traces of denticles (2–3). The LAS structure was not found on genus *Uroctonites* but this was based on limited material for two of the species, mostly adults.

Systematic observations

Table 3 provides a list of non-vaejovid scorpions examined in this study that did *not* exhibit the LAS structure. This list includes representatives of all parvorders, superfamilies, and families of Recent scorpions, spanning over 70 genera. Unlike our analysis of family Vaejovidae, we typically only checked a single species of a genus and not all genera are evaluated, especially in family Buthidae. However, in the families belonging to superfamily Chactoidea, where Vaejovidae resides, we examined almost all genera (except family Supersitioniidae). We can conclude from this examination, though not exhaustive, that the LAS structure in all probability **occurs only in family Vaejovidae**.

We consider this result important since the LAS structure may be a synapomorphy for the family. This conclusion cannot be completely realized until additional material is examined in the *Vaejovis* “nigrescens” and “mexicanus” groups as well as in genera *Pseudouroctonus* and *Uroctonites*. Although we see absence of the LAS in genus *Uroctonites*, additional material must be examined, especially subadults and juveniles. It is clear based on the evidence discussed throughout this paper that the LAS structure does not appear randomly within a species, a species group, or a genus. However, as the statistical analysis also shows, it is highly variable in its number of denticles, and therefore in the length of a denticle row. Due to its very small size, LAS denticles are probably vulnerable to damage or complete destruction due to the scorpion’s stinging mechanism, although we know nothing about the LAS function itself. This would explain why it is more observable in juveniles in some of the groups discussed above.

It might also be observed that the vaejovid genera and *Vaejovis* groups where an absence of this structure

has been noted (we exclude *Paruroctonus hirsutipes* and *Vaejovis occidentalis* from this discussion since they involve only a single specimen in a genus or group where all other species examined exhibited LAS), all share many important taxonomic characters in family Vaejovidae. The “mexicanus” and “nigrescens” groups, in particular, are certainly closely related, the latter’s delineation probably more a product of its lithophilic adaptation than important evolutionary derivations, share the following characters: Chela trichobothria *ib-it* are located basally on the fixed finger, proximal of the last inner denticle (*ID*); chelal palm trichobothrium *Dt* is located proximal of palm midpoint; the genital operculum of the female is separated on the posterior one-fifth; the leg tarsus is equipped with a single pair of ventral distal spinule pairs (with some exceptions in the “mexicanus” group); carapace anterior edge with well developed widely formed indentation; the terminus of the hemispermatophore sperm plug is smooth (after Stockwell, 1989). Genera *Pseudouroctonus* and *Uroctonites* are very closely related, the latter only being diagnosed by “heavier developed” setal pairs on the ventral aspect of leg tarsus, clearly a “character of degree” and probably dubious at best. These two genera share the same characters as the previous two *Vaejovis* groups just discussed, except their leg tarsus is equipped with 2–4 pairs of ventral distal pairs of setae. In addition these two genera have a reduced pectinal tooth count and the ventral median (*V2*) carina of the chela is essentially obsolete.

It is clear from the data presented in this paper that the LAS structure has evolved basally or at least near basally in the family Vaejovidae. Of course, depending on the “final” results of further analysis of the genera and groups discussed above and where the species and/or genera lacking these denticles occur in the topology of the family, will determine whether the LAS structure is a synapomorphy for the family. However, having said this, and based on preliminary cladistic analysis, genus *Uroctonites* does not appear to be a basal taxon in the family and therefore the LAS structure in all likelihood is a synapomorphy for family Vaejovidae.

Other aculear structures

It seems relevant to note here that the basal aculear area, where LAS reside in Vaejovidae, occasionally bears other enigmatic but distinctive, macroscopic structures in scorpions, specifically in the superfamily Chactoidea. One of such structures is an “annular ring” in *Troglocormus*, *Euscorpiops*, and *Alloscorpiops* (Euscorpiidae), found both in males and females (Soleglad & Sissom, 2001, p. 67, figs. 189-191), most prominent in *Alloscorpiops*. Another basal aculear structure is a spectacular inflated “bulb” present in the males of *Anuroctonus* (Chactidae) (Soleglad & Fet,





2004, fig. 20). The functional roles of these structures are not known.

Acknowledgments





We thank Hoang Ngoc Anh, Joe Barnes, Juan Barrios, Matt Braunwalder, Philip Brownell, Matthew Graham, Douglas Gaffin, Dietmar Huber, František Kovařík, Matjaž Kuntner, Wilson Lourenço, Graeme Lowe, Jordi Nebot, Gary Polis, Carles Ribera, Michael Rose, Boris Sket, Rolando Teruel, and Michael Warburg for the generous gift of specimens. We are grateful to Elizabeth Fet for a special gift of scorpions from Wyoming. We thank Janet Beccaloni, Jonathan Coddington, Jürgen Gruber, Blaine Hébert, František Kovařík, Graeme Lowe, Peter Schwendinger, Petra Sierwald, and W. David Sissom for the loan of specimens. This study was supported by Marshall University's Department of Biological Sciences and Department of Chemistry. We are especially grateful to David Neff and Michael Norton for their invaluable help and support of our SEM studies. We also extend our gratitude to František Kovařík and Matthew Graham for reviewing this paper.

References

- FET, V., M. S. BREWER, M. E. SOLEGLAD & D. P. A. NEFF. 2006a. Constellation array: a new sensory structure in scorpions (Arachnida: Scorpiones). *Boletín de la Sociedad Entomológica Aragonesa*, 38: 269–278.
- FET, V. & M. E. SOLEGLAD. 2005. Contributions to scorpion systematics. I. On recent changes in high-level taxonomy. *Euscorpius*, 31: 1–13.
- FET, V., M. E. SOLEGLAD, M. S. BREWER, D. P. A. NEFF & M. L. NORTON. 2006b. Constellation array in scorpion genera *Paruroctonus*, *Smeringurus*, *Vejovoidus*, and *Paravaejovis* (Scorpiones: Vaejovidae). *Euscorpius*, 41: 1–15.
- (PAVLOVSKY, E. N.) PAWLOWSKY, E. N. 1913. Skorpiotomische Mitteilungen. I. Ein Beitrag zur Morphologie der Giftdrüsen der Skorpione. *Zeitschrift für Wissenschaftliche Zoologie*, 105: 157–177.
- SISSOM, W. D. 1990. Systematics, biogeography and paleontology. Pp. 64–160 in G. A. Polis (ed.), *Biology of Scorpions*. Stanford, California: Stanford University Press.
- SISSOM, W. D. 2000. Family Vaejovidae Thorell, 1876. Pp. 503–552 in Fet, V., W. D. Sissom, G. Lowe & M. E. Braunwalder. *Catalog of the Scorpions of the World (1758–1998)*. New York, NY: New York Entomological Society, 690 pp.
- SOLEGLAD, M. E. & V. FET. 2004. The systematics of scorpion subfamily Uroctoninae (Scorpiones: Chactidae). *Revista Ibérica de Aracnología*, 10: 81–128.
- SOLEGLAD, M. E. & V. FET. 2006. Contributions to scorpion systematics. II. Stahnkeini, a new tribe in scorpion family Vaejovidae (Scorpiones: Chactidae). *Euscorpius*, 40: 1–32.
- SOLEGLAD, M. E. & W. D. SISSOM. 2001. Phylogeny of the family Euscorpiidae Laurie, 1896: a major revision. Pp. 25–111 in Fet, V. & P. A. Selden (eds.). *Scorpions 2001. In memoriam Gary A. Polis*. Burnham Beeches, Bucks: British Arachnological Society.
- STOCKWELL, S. A. 1989. *Revision of the Phylogeny and Higher Classification of Scorpions (Chelicerata)*. Ph.D. Thesis, University of Berkeley, Berkeley, California. 319 pp. (unpublished). University Microfilms International, Ann Arbor, Michigan.
- WILLIAMS, S. C. 1970. A systematic revision of the giant hairy scorpion genus *Hadrurus*. *Occasional Papers of the California Academy of Sciences*, 87: 1–62.



Figures 6–9: Laterobasal Aculear Serrations (LAS), ventral view of aculeus base. **6.** *Paravaejovis pumilis*, male, Ciudad Constitución, Baja California, Mexico. **7.** *Vejovoidus longtonguis*, female, Viscaíno Desert, Baja California, Mexico. **8.** *Paruroctonus boreus*, male, Worland, Wyoming, USA. **9.** *Simeringurus mesaensis*, female, Palo Verde Wash, ABDSP, California, USA.



Figures 10–13: Laterobasal Aculear Serrations (LAS), ventral view of aculeus base. **10.** *Paruroctonus arnaudi*, male, El Socorro, Baja California, Mexico. **11.** *Paruroctonus ventosus*, female, El Socorro, Baja California, Mexico. **12.** *Paruroctonus surensis*, male, Las Bombas, Baja California Sur, Mexico. **13.** *Paruroctonus luteolus*, female, Palo Verde Wash, ABDSP, California, USA.