


Euscorpius

Occasional Publications in Scorpiology



Scorpions 2011

John L. Cloudsley-Thompson 90th Birthday Commemorative Volume

Polymorphism and Hybridization in Species of *Hottentotta* Birula, 1908 (Scorpiones: Buthidae)

**Wilson R. Lourenço, Eric Ythier, Mark Stockmann
& John L. Cloudsley-Thompson**

August 2011 — No. 124

Euscorpius

Occasional Publications in Scorpiology

EDITOR: Victor Fet, Marshall University, 'fet@marshall.edu'

ASSOCIATE EDITOR: Michael E. Soleglad, 'soleglad@la.znet.com'

Euscorpius is the first research publication completely devoted to scorpions (Arachnida: Scorpiones). *Euscorpius* takes advantage of the rapidly evolving medium of quick online publication, at the same time maintaining high research standards for the burgeoning field of scorpion science (scorpiology). *Euscorpius* is an expedient and viable medium for the publication of serious papers in scorpiology, including (but not limited to): systematics, evolution, ecology, biogeography, and general biology of scorpions. Review papers, descriptions of new taxa, faunistic surveys, lists of museum collections, and book reviews are welcome.

Derivatio Nominis

The name *Euscorpius* Thorell, 1876 refers to the most common genus of scorpions in the Mediterranean region and southern Europe (family Euscorpiidae).

Euscorpius is located on Website '<http://www.science.marshall.edu/fet/euscorpius/>' at Marshall University, Huntington, WV 25755-2510, USA.

The International Code of Zoological Nomenclature (ICZN, 4th Edition, 1999) does not accept online texts as published work (Article 9.8); however, it accepts CD-ROM publications (Article 8). *Euscorpius* is produced in two *identical* versions: online (ISSN 1536-9307) and CD-ROM (ISSN 1536-9293). Only copies distributed on a CD-ROM from *Euscorpius* are considered published work in compliance with the ICZN, i.e. for the purposes of new names and new nomenclatural acts. All *Euscorpius* publications are distributed on a CD-ROM medium to the following museums/libraries:

- **ZR**, Zoological Record, York, UK
- **LC**, Library of Congress, Washington, DC, USA
- **USNM**, United States National Museum of Natural History (Smithsonian Institution), Washington, DC, USA
- **AMNH**, American Museum of Natural History, New York, USA
- **CAS**, California Academy of Sciences, San Francisco, USA
- **FMNH**, Field Museum of Natural History, Chicago, USA
- **MCZ**, Museum of Comparative Zoology, Cambridge, Massachusetts, USA
- **MNHN**, Museum National d'Histoire Naturelle, Paris, France
- **NMW**, Naturhistorisches Museum Wien, Vienna, Austria
- **BMNH**, British Museum of Natural History, London, England, UK
- **MZUC**, Museo Zoologico "La Specola" dell'Universita de Firenze, Florence, Italy
- **ZISP**, Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia
- **WAM**, Western Australian Museum, Perth, Australia
- **NTNU**, Norwegian University of Science and Technology, Trondheim, Norway
- **OUMNH**, Oxford University Museum of Natural History, Oxford, UK
- **NEV**, Library Netherlands Entomological Society, Amsterdam, Netherlands

Publication date: 7 August 2011

Polymorphism and hybridization in species of *Hottentotta* Birula, 1908 (Scorpiones: Buthidae)

Wilson R. Lourenço¹, Eric Ythier², Mark Stockmann³ & John L. Cloudsley-Thompson⁴

¹ Muséum national d'Histoire naturelle, Département Systématique et Evolution, UMR7205, CP 053, 57 rue Cuvier 75005 Paris, France; e-mail: arachne@mnhn.fr

² SynTech Research, 613 route du Bois de Loyse, 71570 La Chapelle de Guinchay, France, e-mail: eythier@syntechresearch.com

³ Poststraße 69, 49477 Ibbenbüren, Germany

⁴ 10 Battishill Street, Islington, London N1 1TE, United Kingdom

Summary

A new and well documented case of hybridization among scorpions is presented. It was obtained under laboratory conditions between *Hottentotta jayakari* (Pocock) and *Hottentotta salei* (Vachon) specimens of which had been collected in the northern and southern regions of Oman. Hybrids were successfully produced not only from F_0 males and females, but also from F_1 males and females, thereby attesting to the fact that the first generation obtained (F_1), was completely fertile. Both F_1 and F_2 broods were composed of dark and pale morphs, indicating that the juveniles could inherit either one or the other parental phenotype. This report brings new evidence about the true genetic relationship between these two “species”, suggesting that they may correspond only to “morphs” (=phenotypes) of a single polymorphic species.

Introduction

Very few examples of hybridization between different species of scorpions have been reported. Polis & Sissom (1990) dedicate a short section to this question in their chapter on “life history” of scorpions. These authors defined the reported cases as “mistakes” which occasionally occur in the identification of mates, and listed from the literature seven pairs of different species that engage in “promenades” and courtship behaviour. These included four pairs of species from different genera (Auber, 1963; Matthiesen, 1968; Probst, 1972; Le Pape & Goyffon, 1975). Spermatophore deposition and presumably sperm uptake was only observed, however, between congeneric species of *Euscorpius* Thorell (Auber, 1963) and *Androctonus* Ehrenberg (Le Pape & Goyffon, 1975). In the case of mating between a male of *Androctonus australis* (Linnaeus) and a female *Androctonus mauritanicus* (Pocock), 42 supposedly hybrid young were produced.

Several of these observations, however, were extremely empirical and provide no valid evidence of sperm uptake and consequent fecundation (Auber, 1963; Matthiesen, 1968; Probst, 1972). Even in the case of mating between two *Androctonus* species, effective fecundation was not demonstrated (Le Pape & Goyffon, 1975). We will return to this point in the discussion.

More recently, three cases of hybridization between female *Centruroides gracilis* (Latreille) from Mexico and male *Centruroides margaritatus* (Gervais) from Colombia have been observed (Lourenço, 1991). These results suggested strongly that the two species of *Centruroides* may constitute only “morphs” of a widespread polymorphic species. However, since all the offspring were lost before reaching the 3th instar, no elements of the second generation (F_1) could be crossed. Consequently, little could be stated about the true inter-specific relationships between these two species (or ‘morphs’).

In this note we present a new and better documented case of hybridization between two “species” of the genus *Hottentotta*. In this case, “hybrids” were successfully produced not only from F_0 male and female, but also from F_1 male and female, attesting thereby that the first generation (F_1) was completely fertile.

Material Used in the Observations

Scorpions of the species *Hottentotta jayakari* (Pocock, 1895) and *Hottentotta salei* (Vachon, 1980) were received by E.Y. The specimens had been collected respectively in the northern and southern regions of Oman. These are large species which may reach 75–80 mm in total length.



Figures 1–6: 1. F_0 female of *Hottentotta salei* (pale morph). 2. F_0 male of *Hottentotta jayakari* (dark morph). 3. Courtship and mating between F_0 male (dark morph) and F_0 female (pale morph). 4. F_1 brood, on instar I. 5. F_1 brood, on instar II. 6. F_1 adult female (dark morph).

Hottentotta jayakari was described by Pocock (1895), as *Buthus jayakari*, from Muscat in the northern range of Oman. Subsequently, Vachon (1980) described a subspecies of *H. jayakari*, as *Buthotus jayakari salei*, from the Province of Dhofar in the South of Oman. The

diagnostic characters defined by Vachon (1980) were mainly based on the patterns of pigmentation presented by the two populations. In a recent revision of the genus *Hottentotta* by Kovařík (2007), *H. jayakari salei* was raised to the rank of species. Kovařík (2007) argued as



Figures 7–11: 7. F_1 adult male (pale morph). 8. Courtship and mating between F_1 male (pale morph) and F_1 female (dark morph). 9. F_2 brood, on instar II. 10. F_2 juvenile of second instar (dark morph). 11. F_2 juvenile of second instar (pale morph).

follows: “This species was originally described as a subspecies of *H. jayakari*, however the distribution of the two taxons overlap and the species are easily sep-

arated by color” (An overlapping distribution with a zone of parapatry does not exclude the possibility of speciation). Even more recently, in a paper about the

Hottentotta of Oman, Lowe (2010) confirmed the validity of both *H. jayakari* and *H. salei* as distinct species.

Methods

The scorpions were reared by standard methods in plastic terraria of different sizes. These contained layers of soil and sand, 2–3 cm in depth, as well as a few pieces of bark and a small Petri dish containing water. Food, consisting of *Acheta domestica* L. and *Tenebrio molitor* L. larvae, was provided once every 7 to 10 days. Temperatures ranged from 30 to 32°C during the day and 22 to 24°C at night. Humidity was around at 40–50% (R.H.).

Laboratory Observations

Males and females of both species were collected alive, brought to the laboratory and maintained as described above. The courtship and mating behavior of one pair of scorpions was observed on the 30 June 2006. This F_0 pair consisted of one male *H. jayakari* (dark morph) and one female *H. salei* (pale morph) (Figs. 1–3). The female gave birth on 15 July 2007, to an F_1 brood composed of 25 neonates.

Embryonic development in this female lasted for 380 days. After they were carried on their mother's back for 6 days, the first molt of the young scorpions took place on 20 July 2007 (Figs. 4–5). Juveniles began to leave their mother's back at the age of 8 days. Subsequent molts took place at differing ages. The F_1 brood was composed of both dark and pale morphs, attesting to the fact that the juveniles could inherit one or other parental phenotypes (Figs. 6–7).

Five individuals of the F_1 generation, one male (pale morph) and four females (dark morph), were selected for further experiment. The male reached adulthood at the sixth molt (7th instar) after 783 days; the females at the seventh molt (8th instar) at ages varying from 611 to 783 days.

The courtship and mating behaviour of another pair of these scorpions was observed on the 30 November 2009 (Fig. 8). This F_1 pair was composed of a male (pale morph) and a female (dark morph). This female gave birth on 7 July 2010, to an F_2 brood consisting of 18 neonates (Fig. 9).

Embryonic development in this female lasted for 220 days. After they were carried on their mother's back for 5 days, the first molt of the offspring scorpions took place on 11 July 2010. Juveniles began to leave their mother's back at the age of 8 days. Subsequent molts took place, but most of these juveniles are still under observation. The F_2 brood was composed of both dark and pale morphs, attesting once again that juveniles could inherit one of two parental phenotypes (Figs. 10–11).

Discussion

Previous observations of hybridization reported by Auber (1963), Matthiesen (1968) and Probst (1972) were poorly documented, exclusively empirical and did not provided any evidence of sperm uptake or fecundation. In the case, reported by Le Pape & Goyffon (1975), of mating between two *Androctonus* species, effective fecundation was not demonstrated. The two *Androctonus* species used in the experiment occupy totally distinct ranges of distribution (Lourenço, 2005). Le Pape & Goyffon (1975) stated that their chromosomal patterns were totally distinct: 16 for *A. australis* and 24 for *A. mauritanicus*. The suggestion of parthenogenetic reproduction activated by the male spermatozoa is irrelevant because this process has never been observed in the studied cases of parthenogenetic reproduction in scorpions (Lourenço, 2008).

The reported cases of hybridization between *Centruroides gracilis* and *Centruroides margaritatus* (Lourenço, 1991), suggest that these two “species” might only represent “morphs” of a widespread polymorphic species. However, since no members of the F_1 generation could be crossed, it is impossible to determine the relationship between the two species.

The example of hybridization obtained between *Hottentotta jayakari* and *Hottentotta salei* suggests the existence of distinct “morphs” of a polymorphic species:

(a). Hybrids were successfully produced not only between members of F_0 , but also between those of F_1 , which demonstrated that the first generation obtained (F_1), was globally fertile;

(b) F_1 and F_2 broods were composed of dark and pale morphs, showing that the juveniles could inherit one or other parental phenotypes.

Hybridization can take place between closely related subpopulations under laboratory conditions since ecological barriers that separate them in nature do not exist in the laboratory. The different “morphs” observed in nature are very close to one another genetically and, polymorphism can be observed among scorpions in which temporary reproductive isolation does not give rise to genetic incompatibility. However, some minor morphological changes may take place during temporary isolation and, when the subpopulations disperse again the observed variability may no longer be correlated geographically. Such a biogeographical pattern seems to correspond well with certain “complex” populations observed in buthid scorpions (Williams, 1980; Lourenço, 1986, 1988). The species concerned show high vagility, dispersal capacity, ecological plasticity and, in general, have a large geographical distribution. Examples are well known in the genera *Tityus* Koch, *Centruroides* Marx, and could certainly be found among species of the genus *Hottentotta* (Lourenço, 1992).

References

- AUBER, M. 1963. Reproduction et croissance de *Buthus occitanus* Amx. *Annales des sciences naturelles: Zoologie*, 5(2): 273–285.
- KOVAŘIK, F. 2007. A revision of the genus *Hottentotta* Birula, 1908, with description of four new species (Scorpiones, Buthidae). *Euscorpius*, 58: 1–107.
- LE PAPE, G. & M. GOYFFON. 1975. Accouplement interspécifique suivi de parturition dans le genre *Androctonus* (Scorpionida, Buthidae). *Comptes Rendus de l'Académie des Sciences, Paris, Sér. D.*, 280: 2005–2008.
- LOURENÇO, W. R. 1986. Les modèles de distribution géographique de quelques groupes de scorpions néotropicaux. *Comptes Rendu des Séances de la Société de Biogéographie*, 62(2): 61–83.
- LOURENÇO, W. R. 1988. Diversité biologique et modalités de la spéciation chez les scorpions amazoniens; *Tityus silvestris* Pocock, un cas particulier de polymorphisme. *Comptes Rendus de l'Académie des Sciences, Paris*, 306, ser., 3: 463–466.
- LOURENÇO, W. R. 1991. Interspecific hybridation of laboratory reared *Centruroides gracilis* and *Centruroides margaritatus* (Chelicerata, Scorpiones). *Studies on Neotropical Fauna and Environment*, 26 (1): 29–32.
- LOURENÇO, W. R. 1992. Biogéographie évolutive, écologie et les stratégies biodémographiques chez les scorpions néotropicaux. *Comptes Rendu des Séances de la Société de Biogéographie*, 67(4): 171–190.
- LOURENÇO, W. R. 2005. Nouvelles considérations taxonomiques sur les espèces du genre *Androctonus* Ehrenberg, 1828 et description de deux nouvelles espèces (Scorpiones, Buthidae). *Revue suisse de Zoologie*, 112(1): 145–171.
- LOURENÇO, W. R. 2008. Parthenogenesis in scorpions: Some history – new data. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 14(1): 19–44.
- LOWE, G. 2010. Two new species of *Hottentotta* Birula, 1908 (Scorpiones: Buthidae) from northern Oman. *Euscorpius*, 103: 1–23.
- MATTHIESEN, F. A. 1968. On the sexual behaviour of some Brazilian scorpions. *Revista Brasileira de Pesquisas Médicas e Biológicas*, 1(2): 93–96.
- POCOCK, R. I. 1895. On the Arachnida and Myriopoda obtained by Dr. Anderson's collector during Mr T. Brent's expedition to the Hadramaut, South Arabia, with a supplement upon the scorpions obtained by Dr. Anderson in Egypt and the Eastern Soudan. *Journal of the Linnean Society*, 25: 292–316.
- POLIS, G. A. & W. D. SISSOM. 1990. Life history. Pp. 161–223, In: G. A. Polis (ed.). *Biology of Scorpions*. Stanford: Stanford University Press.
- PROBST, P. J. 1972. Zur Fortpflanzungsbiologie und zur Entwicklung der Giftdrüsen beim Skorpion *Isometrus maculatus* (De Geer, 1778) (Scorpiones: Buthidae). *Acta Tropica*, 29(1): 1–87.
- VACHON, M. 1980. Scorpions du Dhofar. The scientific results of the Oman flora and fauna survey 1977 (Dhofar). Scorpions du Dhofar. *Journal of the Oman Studies, Special Report*, 2: 251–263.
- WILLIAMS, S. C. 1980. Scorpions of Baja California, Mexico and adjacent islands. *Occasional Papers of the California Academy of Sciences*, 135: 1–127.